


# Understanding

## ▶ borderline personality disorder



'I wish I had never been diagnosed with BPD. With another diagnosis yet similar behaviour I was treated so differently. Possibly the most painful part of this illness (I will call it that) is the discrimination. And the only reason for this is the diagnosis, not the way I feel, behave or speak, because that was the same before.'

**Borderline personality disorder (BPD) is a controversial diagnosis. This booklet aims to help people to understand when the diagnosis might be given and its consequences. It suggests sources of help for those diagnosed with this problem, their friends and relatives.**



## **What is borderline personality disorder?**

BPD is one of many personality disorders listed in the manuals used by clinicians when they are giving someone a psychiatric diagnosis. The word 'personality' refers to the on-going pattern of thoughts, feelings and outward behaviour that makes us the people we are.

A personality disorder may be diagnosed when it's felt that several areas of someone's personality are causing them or others problems in everyday life. This diagnosis is very controversial, because it implies that someone's whole personality is flawed – rather than just one aspect of them. Some psychiatrists argue that it's impossible to treat someone's personality and that it's wrong to apply medical terms and treatments to a personality. For this reason, it is usually the symptoms of BPD that are addressed in treatment rather than the disorder as a whole. (See Mind's booklet *Understanding Personality Disorders* for more information about this particular group of diagnoses).

Some argue that the term 'borderline' is misleading. Originally, the term was applied to people who seemed to be on the border of being given a diagnosis of schizophrenia. However, now BPD is seen as distinct from schizophrenia diagnoses. The 'borderline' aspect is seen to express being on the border of psychosis. If someone has a psychosis, it means they have beliefs or experiences not shared by others. Those diagnosed with BPD may have these at times of stress.

It has been estimated that three-quarters of those given this diagnosis are women. In the USA, BPD is thought to affect two out of every 100 people. Unfortunately there are no equivalent UK statistics at present. It's a condition that isn't usually diagnosed until adulthood, because the personality is seen as still developing until then.

Because of the controversy surrounding this diagnosis, services are often not readily available. However, there are routes you may be able to take, which are listed later in this booklet.

## **How would a clinician make this diagnosis?**

There are no biochemical or physical tests to tell whether someone does or doesn't have BPD. Instead, clinicians making a diagnosis look to see whether you have had five or more of the following signs, which have been present for at least a year:

- Self-harm and/or repeated attempts or expressions of the desire to commit suicide. An example of this would be cutting yourself. This behaviour can only be counted as one of the criteria for diagnosis; it can't be counted again as demonstrating any of the other symptoms. See Mind's booklet *Understanding Self-harm*, for more information. (Details of this and other booklets mentioned here may be found under *Further Reading* on p. 14.)
- Frantic efforts to avoid being alone due to an intense fear of being abandoned. Others may not see this fear as justified, but you may go to great lengths to avoid being alone. For example, you may say that you will harm someone if they leave.
- Relationship problems where you may see the person you love as absolutely wonderful, able to do no wrong one moment, and then wholly bad the next. Some 'idealisation' is often present in relationships but, here, there will be a pattern of relationships being particularly unsteady and intense.
- A very uncertain, shaky self-image or sense of self. You may feel good whilst you feel loved by someone you think is wonderful. If you later see them as bad, your own sense of self could be affected. You may also have doubts about your sexual identity.

- Two or more areas of your life where your behaviour could cause you harm and be seen as impulsive. Examples would be: spending money extravagantly and having huge debts, having unprotected sex, abusing drugs or alcohol, driving without due care or binge-eating. You may do these things because you're trying to deal with awful feelings of pain or emptiness. (See Mind's booklet *Understanding Eating Distress*.)
- You may have moods that are very difficult to come out of. For example, you may go through long periods – usually a few hours – of extreme irritability, restlessness, unhappiness or anxiety.
- Terrible feelings of emptiness.
- Anger that's inappropriate and intense or difficult to control. You may lose your temper a great deal, experience constant anger or be involved in physical fights. You may feel particularly angry when you think you're being criticised. It's a fine line between assertiveness and the appropriate expression of anger. Anger is often a very difficult feeling for people to acknowledge and deal with but may cause particular problems in the life of someone diagnosed with BPD. (See *How to Deal with Anger*.)
- Periods of paranoia or feeling unreal, when under stress. This might be accompanied by an almost complete lack of physical sensation. At difficult times, you may experience yourself as having more than one personality or feel you are in a trance-like state.

As a result of confusion about your personal identity and a terror of being left alone, you may find yourself clinging to very damaging relationships. Many people who meet the criteria for BPD also meet the criteria for histrionic, narcissistic or antisocial personality disorder. (See *Understanding Personality Disorders*.)

Unfortunately those diagnosed with BPD have a greater risk of committing suicide than the general population. Long-term US studies suggest as many as 9 per cent of those diagnosed with BPD commit suicide. If you are diagnosed with BPD, it's important to know where to turn to if you are feeling suicidal (see *Useful Organisations* on p. 13).

Whilst some people may see themselves in the symptoms of BPD and feel relieved to have a label to apply to the problems they experience, others may be devastated at the idea that their personality is disordered.

It's worth remembering that aspects of almost any type of personality can be found within the pages of the diagnostic manuals. What matters is that you get the help you feel you need. If you feel you have BPD from reading this booklet, be wary of making a self-diagnosis; talk to someone who is medically qualified.

### **What if they've made a mistake in my diagnosis?**

Strictly speaking, a medical diagnosis can only be given by somebody who has been medically trained, a GP or psychiatrist. However, because the term 'borderline' comes from psychoanalytic thinking, you may have this term applied to you by someone who has not been medically trained. There is a recognised and very worrying danger of mistaken diagnosis. Mental health professionals sometimes fall into the trap of applying it to people they have difficulties dealing with, perhaps because of a conflict of personalities. Within the NHS, you are entitled to ask for a second opinion, although this doesn't necessarily mean that your request will be granted.

If you feel your GP or psychiatrist has misunderstood you, and you are having problems getting the help you need, you may find an advocate useful. (For more information about advocates contact the *Mind*infoline or see the *Mind Guide to Advocacy*.)

### **Will I get better?**

It's often thought that personality problems are too deep-seated to be treatable. But this is contradicted by evidence that symptoms may get better as people get older. Some research suggests that after ten years, or so, as many as half of those diagnosed with BPD no longer display enough of the symptoms to deserve the diagnosis. Recent research suggests that talking treatments and medication can reduce the behaviour problems associated with the problem. Day-care programmes may also be useful. (See p. 6 for more information.)



## What causes borderline personality disorder?

### Traumatic experiences

Often, those diagnosed with BPD turn out to have had very traumatic experiences in childhood. You may have experienced the early loss of a parent or be a survivor of childhood sexual and/or physical abuse. You may have been neglected as a child. Such difficult life events are very common in those diagnosed with BPD. (See Mind's factsheet, *Resources for Survivors of Childhood Sexual Abuse*.) The problems associated with BPD may become much worse following a stressful experience, for example, the loss of a loved one or an established routine, such as a job.

### Physical causes

An American psychiatrist, Dr Leland Heller, believes that BPD is a 'neurological illness' probably a form of epilepsy and that it can be managed with appropriate medication and talking treatments. (For more information about his theories, see *Useful Organisations* on p. 13).

## What sort of treatment can I get?

### Talking treatments

Psychotherapy is a relatively long-term talking treatment that aims to find the roots of present feelings and behaviour in your childhood. The relationship you have with the therapist is seen as an important reflection of your past and present relationships. Exploring this relationship can help to break unhelpful patterns of behaviour. The in-depth nature of psychotherapy can make it particularly appropriate for those diagnosed with BPD. Some forms of counselling work in the same way to psychotherapy. Psychodynamic counselling, for example, places great emphasis on childhood experience. (See *Further Reading* on p. 14 and *Useful Organisations* on p. 13 for sources of more information.)

Cognitive behaviour therapy is a more short-term treatment that aims to tackle practical, everyday difficulties with problem-solving techniques. It works towards identifying negative thinking patterns and replacing them with more positive ones.

New therapies have been developed which combine elements of cognitive therapy and psychotherapy. These therapies, for example dialectical behaviour therapy (DBT) and cognitive analytical therapy (CAT), have been found to be particularly useful for people diagnosed with BPD.

People often have high expectations when they enter a talking treatment. It's worth bearing in mind that therapists aren't miracle-workers and that change can take time. If you find it painful to be separated from others, you might want to think about how you will manage breaks in the therapy early on. You could ask when the breaks will occur so that you can look at how you will cope beforehand. There may be times when you think your therapist is wonderful and times when you may hate him or her. It may help you to express these feelings, so that you can look at them together.

If you are interested in pursuing a talking treatment, you could talk to your GP about the possibility of seeing someone through the NHS, or getting treatment subsidised.

### Therapeutic communities

The NHS runs some in-patient therapeutic communities that specialise in treating clients with personality disorders (see *Useful Organisations*, on p. 13). In a therapeutic community, staff and residents share responsibility for tasks and decisions. If you decide to go into a therapeutic community, you will need to be prepared to talk about your life with others before the group decides whether to give you a place. This can be hard, especially if it's the first time you have talked in front of a group in this way. Once part of the community, you would be encouraged to talk about your feelings about others' behaviour in group discussions. This may seem difficult at first but it can be very beneficial. It may give you the opportunity to see how others react to you and what you say. You can then think about what you like and what you want to change about yourself. Some, but not all communities may offer you individual therapy and, possibly, medication.

## Alternative therapies

There are a whole range of alternative therapies, which some people find useful, from acupuncture to yoga. See Mind's booklet *A-Z of Alternative and Complementary Therapies*.



## What about medication?

### Antidepressants

Research has found low levels of the chemical serotonin in people diagnosed with BPD who have committed impulsive acts of violence. Various factors can cause changes to serotonin levels. Some of the antidepressants work to increase levels of serotonin. (For more information about antidepressants and their side-effects see Mind's booklet *Making Sense of Treatments and Drugs: Antidepressants*.)

### Major tranquillizers

Major tranquillizers are also referred to as antipsychotics. These drugs may be prescribed to help with feelings of unreality or paranoia. They should be prescribed with caution as they can have distressing side-effects especially in long-term use. (For more information, see Mind's booklet *Making Sense of Treatments and Drugs: Major Tranquillizers*.)



## What happens if I can't cope?

### Crisis services

In response to demand from mental health service-users, crisis services have been developed in some areas. In these services, the emphasis is on talking treatments and informal support. A crisis service may be somewhere safe to stay, or an out-of-hours telephone helpline. (Mind's factsheet *Crisis Services* gives further details.)

### Hospital

At times of great distress you may feel you need to be somewhere safe. This could mean going into hospital. It can be upsetting to be around others who are in pain, however, and you may feel a lack of privacy and support. Service-user or patient groups based in the hospital can be useful and supportive.

Most admissions are voluntary but if you are regarded as a danger to yourself or to others, but you don't wish to go, you may be admitted compulsorily under the Mental Health Act 1983. Mind's *Rights Guides* (see p. 14) explain your rights with regards to mental health law. The Mind *infodine* can refer you to Mind's Legal Advice line.

## **What should family and friends do?**

It's important not to see someone purely in terms of their diagnosis. People with BPD can have very low self-esteem and it can help them enormously if you can emphasise the positive parts of their personality.

It can be extremely difficult caring for someone with BPD. They may try very hard to control you, because they feel so out of control of themselves. There may be periods when they refuse to talk to you or when they rage at you. This can be very painful and may remind you of times when you felt powerless as a child.

You may find that the person panics and perhaps reacts very angrily when you want to leave or to go somewhere. They may beg you to stay, or hurl words of abuse. It can help if you focus on how they are feeling, rather than trying to argue them out of their fears.

### Looking after yourself

It's very important to look after yourself and to remember that you need time to yourself, if you are to care for others. If you are experiencing problems – for example, if the person calls you many times a day at work – it's vital to set down some boundaries. It might be important to decide how often you will be available. This can be hard to stick to, especially if you are being threatened, and you may need to enlist other people's help. Offer to help draw up a list of numbers the person could call when they feel afraid.

A person diagnosed with BPD may feel that they have no control over their feelings; they may blame you for everything. Make sure you have someone you can turn to, to help you look at what is happening and make sure you don't take the blame for absolutely everything. Nobody deserves to be abused.

There are organisations that can help you talk about the situation and make decisions about what you're going to do. You may need support in the form of a self-help group or some kind of talking treatment (see *Useful Organisations* opposite for more information).

In an emergency

If you feel that the person you care for is a serious danger to themselves or others, you might need to think about the last resort of compulsory admission to hospital. The 'nearest relative' as defined under the Mental Health Act 1983 can request a Mental Health Assessment from a social worker specially trained in mental health law. The social worker would decide, with the help of medical advice, what the treatment options should be and whether the person needs to be detained (see Mind's *Rights Guides*, details on p. 14).

## i

### References

- Amongst Ourselves: A self help guide to living with Dissociative Identity Disorder* T. Alderman, K. Marshall (New Harbinger 1998)
- A Possible New Name for Borderline Personality Disorder* Dr L. M. Heller
- Advice for Carers on BPD* First Steps to Freedom
- Diagnosis* K. Darton (OpenMind 95, Jan/Feb 1999)
- DSMIV-IV, Diagnostic and Statistical Manual of Mental Disorders* (American Psychiatric Association 2000)
- ICD10 Classification of Mental and Behavioural Disorders* (World Health Organisation 1992)
- Making us Crazy* H. Kutchins, S. Kirk (Constable 1999)
- Medical Treatment of the Borderline Personality Disorder* Dr L. M. Heller (1998)
- Personality Disorder: A Way Forward?* H. Castillo, D. Tallis (Mind Annual Conference 2000)
- Should Psychiatrists Treat Personality Disorders?* P. Moran (Maudsley Discussion Paper No. 7)
- The Structure and Development of Borderline Personality Disorder: A proposed model* A. Ryle (British Journal of Psychiatry 170,1997)
- The Care Programme Approach and Risk Assessment of Borderline Personality Disorder* P. Whewell, D. Bonanno (Psychiatric Bulletin 24, 381-384, 2000)

## Useful organisations

Association of Therapeutic Communities  
Pine Street Day Centre  
13-15 Pine Street  
London EC1R 0JH  
tel./fax: 020 8950 9557  
e-mail: [post@therapeuticcommunities.org.uk](mailto:post@therapeuticcommunities.org.uk)  
web: [www.therapeutic.communities.org.uk](http://www.therapeutic.communities.org.uk)  
Produces a directory of therapeutic communities

Borderline UK  
PO Box 42  
Cockermouth  
Cumbria CA13 0WB  
e-mail: [info@borderlineuk.co.uk](mailto:info@borderlineuk.co.uk)  
web: [www.borderlineuk.co.uk](http://www.borderlineuk.co.uk)  
User-led network of people with a BPD diagnosis in the UK.  
Website includes links to on-line support groups

British Association for Behavioural and Cognitive Psychotherapies  
PO Box 9  
Accrington BB5 0XB  
tel: 01254 875277  
fax: 01254 239114  
e-mail: [babcp@babcp.com](mailto:babcp@babcp.com)  
web: [www.babcp.com](http://www.babcp.com)  
Full directory of psychotherapists available. Can be searched on their website by specialism 'personality disorders'

First Steps to Freedom  
7 Avon Court  
School Lane  
Kenilworth  
Warwickshire CV8 2GX  
helpline: 01926 851608  
fax: 0870 164 0567  
e-mail: [info@firststeps.demon.co.uk](mailto:info@firststeps.demon.co.uk)  
web: [www.firststeps.demon.co.uk](http://www.firststeps.demon.co.uk)  
Supports friends and relatives of those with BPD

National Association for People Abused in Childhood (NAPAC)  
Union House c/o BSS  
Shepherds Bush Green  
London W12 8UA  
tel. 020 8735 5009  
fax: 020 8735 5099  
e-mail: [info@napac.org.uk](mailto:info@napac.org.uk)  
web: [www.napac.org.uk](http://www.napac.org.uk)  
Postal support, advice and guidance for adult survivors of any form  
of childhood abuse – sexual, physical or emotional

National Drugs Helpline  
tel. 0800 776600  
minicom: 0800 917 8765  
web: [www.ndh.org.uk](http://www.ndh.org.uk)  
Free 24 hour helpline for information and advice about drug use

The Cassel Hospital  
1 Ham Common  
Richmond  
Surrey TW10 7JF  
tel. 020 8940 8181  
fax: 020 8237 2996  
web: [www.thecasselhospital.org.uk](http://www.thecasselhospital.org.uk)  
e-mail: [khealy.cassel@btinternet.com](mailto:khealy.cassel@btinternet.com)  
In-patient therapeutic community for people with personality  
disorders. Offers individual psychotherapy and medication where  
appropriate

The Henderson Hospital  
2 Homeland Drive  
Sutton  
Surrey SM2 5LT  
tel. 020 8661 1611  
fax: 020 8770 3676  
e-mail: [sgarner@swlstg-tr.nhs.uk](mailto:sgarner@swlstg-tr.nhs.uk)  
web: [www.swlstg-tr.org.uk/henderson/default.htm](http://www.swlstg-tr.org.uk/henderson/default.htm)  
Therapeutic communities for people with personality disorders.  
Relies on group psychotherapy, and 'living and learning' experience

The Samaritans  
helpline: 08457 90 90 90  
e-mail: [jo@samaritans.org](mailto:jo@samaritans.org)  
A 24-hour emergency helpline

UK Council for Psychotherapy (UKCP)  
167-169 Great Portland Street  
London W1N 5PF  
tel. 020 7436 3002  
fax: 020 7436 3013  
e-mail: [ukcp@psychotherapy.org.uk](mailto:ukcp@psychotherapy.org.uk)  
web: [www.psychotherapy.org.uk](http://www.psychotherapy.org.uk)  
Information about properly accredited psychotherapists

YoungMinds  
102-108 Clerkenwell Road  
London EC1M 5SA  
tel. 020 7336 8445  
fax: 020 7336 8446  
parents information service: 0800 018238  
e-mail: [enquiries@youngminds.org.uk](mailto:enquiries@youngminds.org.uk)  
web: [www.youngminds.org.uk](http://www.youngminds.org.uk)  
Information for parents and those concerned about the mental health of a child or adolescent

## **websites**

The Behavioural Technology Transfer Group  
[www.behavioraltech.com](http://www.behavioraltech.com)  
Contains a section on Dialectical Behaviour Therapy

[www.biologicalunhappiness.com](http://www.biologicalunhappiness.com)  
About BPD as a neurological illness

Virtual Institute of Severe Personality Disorder (VISPED)  
[www.doh.gov.uk/hspscb/visped.htm](http://www.doh.gov.uk/hspscb/visped.htm)  
DOH-sponsored research and development project on BPD

## Further reading

- The Anger Control Workbook* M. McKay, P. Rogers (New Harbinger Press 2000) £14.99
- The Assertiveness Workbook* R. Paterson (New Harbinger Press 2000) £12.99
- A-Z of Complementary and Alternative Therapies* (Mind 2000) £3.50
- Beyond Survival: Living well is the best revenge* Y. Dolan (BT Press 2000) £15.50
- Factsheet: *Cognitive Behavioural Therapy* (1999) 50p
- Factsheet: *Crisis Services* (Mind 2001) £1
- Factsheet: *Resources for Survivors of Child Sexual Abuse* (Mind 2000) 50p
- The Hurt Yourself Less Workbook* (NSHN 1998) £25
- How to Cope with Doubts about your Sexual Identity* (Mind 1999) £1  
*Making Sense of Treatments and Drugs:*
- Antidepressants* (Mind 1998) £1
- Major Tranquillizers* (Mind 1999) £1
- Making us Crazy – DSM: The psychiatric bible and the creation of mental disorders* H. Kutchins, S. Kirk (Constable 1999) £14.99
- Managing Anger* G. Lindenfield (Thorsons 1993) £6.99
- The Mind Guide to Advocacy* (Mind 2001) £1  
*Mind Rights Guides:*
- 1. *Civil Admission to Hospital* (Mind 2001) £1
- 2. *Mental Health and the Police* (Mind 1995) £1
- 3. *Consent to Medical Treatment* (Mind 1995) £1
- 4. *Discharge from Hospital* (Mind 1995) £1
- 5. *Mental Health and the Courts* (Mind 1995) £1
- 6. *Supervision Registers and Supervised Discharge* (Mind 1997) £1
- National Self Harm Network Information Pack* (NSHN 1998) £3.50
- Overcoming Low Self-Esteem* M. Fennell (Robinson 1999) £7.99
- Overcoming Traumatic Stress* C. Herbert, A. Wetmore (Robinson 1999) £7.99
- Recovery: An alien concept*, R. Coleman (Handsell Publishing 1999) £10
- Understanding Attention Deficit Hyperactivity Disorder* (Mind 1997) £1
- Understanding Eating Distress* (Mind 2000) £1
- Understanding Paranoia* (Mind 2000) £1
- Understanding Personality Disorders* (Mind 2000) £1
- Understanding Self-harm* (Mind 2000) £1

# order form

For a catalogue of publications from Mind, send an A4 SAE to the address below.

If you would like to order any of the titles listed here, please photocopy or tear out these pages, and indicate in the appropriate boxes the number of each title that you require.

Please add 10% for postage and packing, and enclose a cheque for the whole amount, payable to Mind. Return your completed order form together with your cheque to:

Mind Mail Order  
15-19 Broadway  
London  
E15 4BQ  
tel. 020 8221 9666  
fax: 020 8534 6399  
e-mail: [publications@mind.org.uk](mailto:publications@mind.org.uk)  
web: [www.mind.org.uk](http://www.mind.org.uk)  
(Allow 28 days for delivery)

Please send me the titles marked above. I enclose a cheque (including 10% for p&p) payable to Mind for £

Name
Address
Postcode
tel.

# Mind works for a better life for everyone with experience of mental distress

Mind does this by:

- advancing the views, needs and ambitions of people with experience of mental distress
- promoting inclusion through challenging discrimination
- influencing policy through campaigning and education
- inspiring the development of quality services which reflect expressed need and diversity
- achieving equal civil and legal rights through campaigning and education

The values and principles which underpin Mind's work are:  
autonomy, equality, knowledge, participation and respect

For details of your nearest Mind association and of local services contact Mind's helpline, *MindinfoLine*: 0845 7660 163 Mon – Fri 9.15am – 5.15pm.

For interpretation, *MindinfoLine* has access to 100 languages via Language Line. Typetalk is available for people with hearing or speech problems who have access to a minicom. To make a call via Typetalk dial 0800 959598, fax: 0151 709 8119.

Scottish Association for Mental Health tel. 0141 568 7000

Northern Ireland Association for Mental Health tel. 02890 328474



**This booklet was written by Louise Flory**

ISBN 1-903567-19-X

© Mind 2001

No reproduction without permission

Mind is a registered charity No. 219830

Mind (National Association for Mental Health)

15-19 Broadway, London E15 4BQ

tel. 020 8519 2122, fax: 020 8522 1725

web: [www.mind.org.uk](http://www.mind.org.uk)